

COMPUTING THE WALLS ASSOCIATED TO BRIDGELAND STABILITY CONDITIONS ON PROJECTIVE SURFACES

ANTONY MACIOCIA

ABSTRACT. We derive constraints on the existence of walls for Bridgeland stability conditions for general projective surfaces. We show that in suitable planes of stability conditions the walls are bounded and derive conditions for when the number of walls is globally finite. In examples, we show how to use the explicit conditions to locate walls and sometimes to show that there are no walls at all.

INTRODUCTION

Throughout this paper we let X be a smooth complex projective surface. The notion of stability condition for a triangulated category (and, in particular, for the derived category of coherent sheaves on X) was introduced by Bridgeland (see [Bri07]). Bridgeland went on in [Bri08] to construct explicit families of stability conditions on K3 and abelian surfaces. Arcara and Bertram ([ABL07]) then showed that these examples work for all smooth complex projective surfaces. We shall be exclusively interested in these examples and their associated walls. We will call these **basic** (Bridgeland) stability conditions.

There are various reasons for studying such stability conditions now well documented in the literature. For example, the preservation of stability under Fourier-Mukai transforms correspond to the non-existence of walls. The Göttsche conjecture and Donaldson-Thomas invariants are (conjecturally) related to wall crossing formula in such stability spaces.

One particular use is to help us to compute explicit moduli spaces of objects in the derived category of a variety. At a boundary of the space of stability conditions the stability condition becomes the more familiar one of (twisted) Gieseker stability. We can construct these spaces using techniques of Geometric Invariant Theory and often they are related to other moduli spaces such as moduli spaces of connections satisfying certain differential equations. But for fixed characteristic classes, the space of stability conditions is divided into chambers and once we cross a wall into a different chamber the objects representing points of the corresponding moduli space need not be torsion-free sheaves or indeed be sheaves at all. In order to better understand these moduli spaces (including their existence) our first step is to locate the walls separating the chambers. These walls arise from a combination of numerical constraints and constraints coming from the hom sets of objects in the derived category. Indeed, understanding better which walls can exist should help us understand the derived category and its sets of maps. The numerical

Date: February 22, 2012.

1991 Mathematics Subject Classification. 14F05, 14D20, 14J60, 18E30, 14N35, 14C20.

Key words and phrases. Moduli space, Bridgeland stability, walls.

constraints turn out to be essentially quadratic in suitable coordinates in the stability manifold $\text{Stab}(X)$ and so are (real) quadrics. If we restrict to certain real half-planes in $\text{Stab}(X)$ then these are conic sections with centres on the boundary of the half-planes. We will prove that, provided we choose the plane correctly, these are heavily constrained to be **nested**. By this we mean that the conics lie entirely inside each other so that they can never intersect or lie in two halves of the plane divided by a straight line (see Figure 1 in the text for a typical example). We prove:

Theorem 3.1 (Bertram's Nested Wall Theorem). *Let X be a smooth complex projective surface. Pick an ample $\omega \in \text{NS}(X)$ and a real class $\beta \in \text{NS}_{\mathbb{R}}(X)$. Then for any class $v \in N(X)$ representing the Chern character of a μ -stable sheaf or torsion sheaf on X there is a half-plane $\Pi = \Pi_{\beta, \omega, 1}$ in $\text{Stab}(X)$ which contains the basic Bridgeland stability condition $(\mathcal{A}_{\beta, \omega}, Z_{\beta, \omega})$ such that the intersection of the walls with Π are nested.*

In particular, if the Picard Group $\rho(X) = 1$ the walls in the space of basic stability conditions are nested.

This theorem was conjectured by Aaron Bertram and there was already a lot of evidence for it in the Picard rank 1 case on \mathbb{P}^2 and on abelian surfaces (see [MM11] for the abelian surface case). The key ingredients are the Bogomolov inequality and the Hodge Index Theorem (just as in [ABL07]) applied to v .

The proof uses only the numerical constraints (we will call the resulting walls **pseudo-walls** in what follows) and actually proves the stronger result that the radii of the circles depends only on the position of the centre of the circles. It also follows from the analysis that given a Chern character v there exists a real number C_0 such that every wall must cross the line $s = C_0$. This is particularly helpful to help us locate walls by providing bounds on the Chern classes of destabilizing objects. The number C_0 is given by the formula

$$\frac{c_1(v) \cdot \omega}{r(v)\omega^2} - \sqrt{F},$$

where F is given by Equation 2.1. In the special case of Picard rank 1, $F = \Delta(v)/r(v)^2\omega^2$, where $\Delta(v)$ is the discriminant of v . We show that when $F = 0$ there can be no walls at all. In other words, a Bogomolov critical μ_ω -stable sheaf must always be μ_Z -stable for any basic Bridgeland stability condition (\mathcal{A}, Z) .

Using our constraints on the walls and the fact that walls must correspond to actual destabilising objects we show that the radii of the walls is always bounded above (and we give explicit computable bounds). More precisely, we shall show:

Theorem 3.10. *Fix a Chern character v of a μ_ω -stable sheaf or a torsion sheaf with $F > 0$. In the half-plane $\Pi_{\beta, \omega, u}$ parametrized by (s, t) with $t > 0$, there are real numbers C_0 and R_0 depending on u such that all of the walls are contained in the semi-circle with centre $(C_0, 0)$ and radius R_0 . In particular, the radii of the walls are bounded above by R_0 .*

Together with local finiteness of the walls we can then give a new more direct proof of a result in [LQ11] which shows that for a given β and ω , there is some $M \in \mathbb{R}$ such that there are no mini-walls (these are just the intersection of walls with rays $t > 0$ in our planes)

for $t \in [M, \infty)$. We also show that for all but one ray in our plane the number of walls is actually globally finite.

We finish with a number of examples. Firstly, we show how to use C_0 to quickly show that, for certain Chern classes v , walls cannot exist. We then look again at the [MM11] situation of an irreducible principally polarized abelian surface and rank 1 sheaves. We also consider the same situation but for a product abelian surface and this allows us to construct an example to show that for other choices of plane, the walls do not nest.

1. STABILITY CONDITIONS AND THEIR WALLS

1.1. The numerics of a surface. Let X be a smooth complex projective surface and let $D(X) = D(\text{Coh } X)$ denote its bounded derived category of coherent sheaves. We let $N(X)$ denote the **numerical Grothendieck group** of X . For a coherent sheaf A on X we let $v(A)$ denote its image in A . This can be extended to the derived category by defining $v(a) = \sum_i (-1)^i v(A^i)$, where $a \in D(X)$ and we use the convenient abbreviated notation $A^i = H^i(a)$.

To be more definite we shall identify $N(X)$ with $\mathbb{Z} \oplus \text{NS}(X) \oplus \mathbb{Z}[\frac{1}{2}]$, where the factors correspond to $r(A)$, $c_1(A)$ and $\text{ch}_2(A)[X]$. We shall freely write $v(a) = (r(a), c_1(a), \text{ch}_2(a)[X])$ corresponding to this decomposition and abuse notation by writing $\text{ch}_2(E) = \text{ch}_2(E)[X]$. Note that the rank of $\text{NS}(X)$ is the **Picard number** $\rho(X)$ of X . We denote the intersection pairing on $\text{NS}(X)$ by $(c, d) \mapsto c \cdot d$. Its signature is $(1, \rho(X) - 1)$. This is extended to a pairing of signature $(2, \rho(X))$ on $N(X)$ by $(v, w) \mapsto v_1 w_3 + v_3 w_1 + v_2 \cdot w_2$.

For any \mathbb{Z} -module M we write $N_M(X) = N(X) \otimes M$ and $\text{NS}_M(X) = \text{NS}(X) \otimes M$. In particular, when $M = \mathbb{Q}, \mathbb{R}$ or \mathbb{C} . We abuse notation and write $\alpha \in \text{NS}_M(X)$ for the image of some $\alpha \in \text{NS}_N(X)$, where M is an N -module. Suppose $\omega \in \text{NS}(X)$ is an ample class and given any non-zero element β of $\text{NS}_{\mathbb{R}}(X)$ we write $\beta = b\omega + \gamma$ for the unique element $\gamma = \gamma(\beta)$ of $\text{NS}_{\mathbb{R}}(X)$ which is orthogonal to ω . We let $\tilde{\gamma}$ denote a choice of element of $\text{NS}(X)$ which maps to γ in $\text{NS}_{\mathbb{R}}(X)$. We can play the same game in $\text{NS}_{\mathbb{Q}}(X)$ but in that case there is a canonical choice for $\tilde{\gamma}$ given by $n\gamma$ for the least positive integer n . This trick is also used in Kawatani [Kaw11] to analyse slightly different stability conditions.

For a fixed ample class $\omega \in \text{NS}(X)$ we let $\mu_{\omega}(v) = c_1(v) \cdot \omega / r(v)$ for any vector $v \in N_R(X)$, interpreting it as $\pm\infty$ when $r(v) = 0$. We let $\mu_{\omega}(a) = \mu_{\omega}(v(a))$ for an object $a \in D(X)$. As is well known, any sheaf $E \in \text{Coh}(X)$ has a unique Harder-Narasimhan filtration by μ_{ω} -semistable sheaves $\{E_i\}_{i=1}^n$ with $\mu_{\omega}(E_i) > \mu_{\omega}(E_{i+1})$. We write $\mu_{\omega}^+(E) = \mu_{\omega}(E_1)$ and $\mu_{\omega}^-(E) = \mu_{\omega}(E_n)$. We call n the **length** of the Harder-Narasimhan filtration.

1.2. Distinguished subvarieties of $\text{Stab}(X)$. Now, fix an ample class $\omega \in \text{NS}(X)$ and another class $\beta \in \text{NS}_{\mathbb{R}}(X)$. Then we can define an abelian subcategory $\mathcal{A}_{\beta, \omega}$ of $D(X)$ by the following tilt with respect to $\text{Coh}(X)$:

$$\begin{aligned} T_{\beta, \omega} &= \{E \in \text{Coh}(X) : E \text{ is torsion or } \mu_{\omega}^-(E) > \beta \cdot \omega\} \\ F_{\beta, \omega} &= \{E \in \text{Coh}(X) : E \text{ is torsion-free and } \mu_{\omega}^+(E) \leq \beta \cdot \omega\} \end{aligned}$$

Notice that, for any $t > 0$, $\mathcal{A}_{\beta,t\omega} = \mathcal{A}_{\beta,\omega}$. We can make these abelian subcategories into a Bridgeland stability condition by defining

$$Z_{\beta,\omega}(v) = -\exp(-\beta - i\omega) \cdot v$$

in $N_{\mathbb{C}}(X)$. The details are discussed in [ABL07, §2]¹. We shall refer to these as **basic stability conditions**. To be explicit note that

$$Z_{\beta,\omega}((r, \theta, z)) = -z + \theta \cdot \beta - \frac{r}{2}(\beta^2 - \omega^2) + i(\theta - r\beta) \cdot \omega.$$

We can use this to describe subvarieties of $\text{Stab}(X)$. Firstly $(\mathcal{A}_{\beta,\omega}, Z_{\beta,\omega})$ is a point of $\text{Stab}(X)$ (actually, a non-trivial fact). We can also consider the set $(\mathcal{A}_{\beta,\omega}, Z_{\beta,t\omega})$ as a ray, denoted $\Xi_{\beta,\omega}$, in $\text{Stab}(X)$. Note that this starts at the open boundary of $\text{Stab}(X)$ (corresponding to $t = 0$). If $\beta \neq 0$, we can also consider the “plane” (or more properly, “half-plane”) $\{(\mathcal{A}_{s\beta,\omega}, Z_{s\beta,t\omega}) : s \in \mathbb{R}, t > 0\}$. This turns out to be hard to describe in general and we modify this to a family of planes as follows. Write $\beta = b\omega + \gamma$ as above and then consider the (usually) half 3-space of stability conditions

$$\Omega_{\beta,\omega} = \{(\mathcal{A}_{sw+u\gamma,\omega}, Z_{sw+u\gamma,t\omega}) : s, u \in \mathbb{R}, t > 0\}.$$

We can now consider the u -indexed family of planes given as the subspaces of $\Omega_{\beta,\omega}$ where u is fixed. We shall denote each of these planes by $\Pi_{\beta,\omega,u}$.

- Notes 1.1.** (1) If $\gamma = 0$ (so $\beta \propto \omega$) then $\Omega_{\beta,\omega}$ is just a half-plane $\Pi_{\beta,\omega,0}$. If $\rho(X) = 1$ then γ must be zero.
 (2) The ray $\Xi_{0,\omega}$ is contained in $\Pi_{\omega,\omega,0}$.
 (3) More generally, if $b = \beta \cdot \omega = 0$ but $\beta \neq 0$ then $\Omega_{\beta,\omega} = \Omega_{\beta+\omega,\omega}$ is still three dimensional.
 (4) Suppose $\alpha \in \text{NS}_{\mathbb{Q}}(X)$ is non-zero and orthogonal to both γ and ω . Then $Z_{\beta,\omega}(v) = Z_{\beta,\omega}(v + \alpha)$.

Example 1.2. Consider the case of a product abelian surface $(\mathbb{T}, \omega) = (\mathbb{T}_1 \times \mathbb{T}_2, \ell_1 + \ell_2)$. Assume $\text{NS}(\mathbb{T}) = \langle \ell_1, \ell_2 \rangle$. Then, over \mathbb{Q} we have $\text{NS}_{\mathbb{Q}}(\mathbb{T}) = \langle \omega, \ell_1 - \ell_2 \rangle$. Then $\Omega_{\ell_1 - \ell_2, \omega}$ contains all of the possible basic stability conditions.

1.3. Walls and pseudo-walls. Given a stability condition (\mathcal{A}, Z) , we say that an object $a \in \mathcal{A}$ is **semistable** if for all proper injections $b \rightarrow a$ in \mathcal{A} we have $\phi(b) \leq \phi(a)$, where $\phi(b) = \arg(Z(v(b)))$. Equivalently, we can use the slope function:

$$\begin{aligned} \mu_Z(b) &= -\frac{\text{Re } Z(v(b))}{\text{Im } Z(v(b))} \\ &= \frac{z - \theta \cdot \beta + \frac{r}{2}(\beta^2 - \omega^2)}{(\theta - r\beta) \cdot \omega}, \end{aligned}$$

where $v(b) = (r, \theta, z)$ and $Z = Z_{\beta,\omega}$.

Now fix a non-zero class $v \in N(X)$ and for the following definition we assume that $v = v(a)$ for some object a of $D(X)$.

¹In fact, we do not seem to need the Harder-Narasimhan property in our analysis.

Definition 1.3. We say that a class $w \in N(X) \setminus \langle v \rangle$ is **critical** for v if each of the following condition hold:

There is a $\sigma \in \text{Stab}(X)$ and objects $a, b \in \mathcal{A}_\sigma$ with an injection $b \rightarrow a$ in \mathcal{A}_σ such that $v(a) = v$ and $v(b) = w$ and $\mu_\sigma(a) = \mu_\sigma(b)$.

If w is critical for v then we write $W_w^v \subset \text{Stab}(X)$ for the set of such σ and call it the **wall** corresponding to w . We drop the v from the notation if it is understood.

Notes 1.4. If there is no $a \in D(X)$ such that $v(a) = v$ then there can be no critical w . Generally, we assume that v is the Chern character of some object.

If a wall exists then it must be a real codimension 1 subspace of $\text{Stab}(X)$. The complement of the union of all of the walls is a disjoint union of chambers with the property that if for any σ_0 in a chamber and object a is σ_0 -stable, then it remains σ -stable for all σ in the same chamber. One would hope that there is a good coarse moduli scheme for such objects and crossing a wall would correspond to a surgery on these moduli spaces. We do not pursue the details here but instead use this as motivation to provide information about finding such walls. To help us do this we introduce a weaker notion of **pseudo-wall**.

Definition 1.5. We say that a class $w \in N(X) \setminus \langle v \rangle$ (respectively, $w \in N_{\mathbb{R}}(X) \setminus \langle v \rangle$) is **\mathbb{Z} -critical** (respectively, **\mathbb{R} -critical**) for v if there is a stability condition σ for v such that $\mu_Z(w) = \mu_Z(v)$.

Remark 1.6. It is clear that a wall is also an integral pseudo-wall and an integral pseudo-wall is also a real pseudo-wall. But the converses are generally false.

Remark 1.7. The set of pseudo-walls (real or integral) do not depend very much on X .

Remark 1.8. Note that we will be interested in these purely numerical constraints on the existence of walls (embodied in the pseudo-walls) which are well defined for any $v \in N_M(X)$ but in the applications we will always want v to be a valid class corresponding to either a μ_ω -semistable sheaf E or a torsion sheaf E . The Bogomolov inequality then implies that $c_1(v)^2 \geq 2r(v) \text{ch}_2(v)$. Note also that if $v = v(E)$ then $E \in T_{s\omega+u\gamma,\omega}$ and so either $r(E) = 0$ or $\mu_\omega(E) > s\omega^2$.

Definition 1.9. We say that $v \in N_M(X)$ (for any M) is a **Bogomolov class** if it satisfies the Bogomolov inequality:

$$c_1(v)^2 \geq 2r(v) \text{ch}_2(v).$$

Remark 1.10. Finally, in this section observe that twisting v by L shifts the walls and pseudo-walls by a unit to the right.

2. PURELY NUMERICAL CONSTRAINTS

Now fix a Bogomolov class $v = (x, \theta, z)$ and an ample class ω in $\text{NS}(X)$. Assume for the moment that $\rho(X) > 1$ and fix a class $\gamma \in \text{NS}_{\mathbb{Q}}(X)$ which is orthogonal to ω . Write $\text{NS}_{\mathbb{Q}}(X) = \langle \omega \rangle \oplus \langle \gamma \rangle \oplus \langle \omega, \gamma \rangle^\perp$. Decompose $\theta = y_1\omega + y_2\gamma + \alpha$, where $\alpha \in \langle \omega, \gamma \rangle^\perp$. If $\rho(X) = 1$ then we can usually just let $\gamma = 0$. Let $Z = Z_{s\omega+u\gamma,t\omega}$. We shall also abbreviate

$\omega^2 = g$ and $\gamma^2 = -d$. Note that $d \geq 0$ by the Hodge Index Theorem and $d = 0$ if and only if $\gamma = 0$. Then

$$\mu_Z(v) = \frac{z - sy_1g + uy_2d + \frac{x}{2}(s^2g - u^2d - t^2g)}{(y_1 - xs)gt}.$$

Let $w = (r, c_1\omega + c_2\gamma + \alpha', \chi)$. Then

$$\begin{aligned} \mu_Z(w) - \mu_Z(v) = & \frac{(y_1 - xs)\left(\chi - sc_1g + uc_2d + \frac{r}{2}(s^2g - u^2d - t^2g)\right) \\ & - (c_1 - rs)\left(z - sy_1g + uy_2d + \frac{x}{2}(s^2g - u^2d - t^2g)\right)}{(c_1 - rs)(y_1 - xs)t} \end{aligned}$$

The numerator $\times 2$ then equals (collecting terms in s and t):

$$\begin{aligned} & g(xc_1 - y_1r)s^2 - 2(\chi x - rz + c_2udx - y_2udr)s + \\ & g(xc_1 - y_1r)t^2 - 2zc_1 + 2c_2udy_1 + xu^2dc_1 - 2y_2udc_1 + 2\chi y_1 - ru^2dy_1. \end{aligned}$$

Now assume $\mu_\omega(v) \neq \mu_\omega(w)$ (so we can divide by $c_1x - ry_1$). Completing the squares we can write this as

$$g(xc_1 - ry_1)((s - C)^2 + t^2 - D - C^2),$$

where

$$\begin{aligned} C &= \frac{x\chi - rz + ud(xc_2 - ry_2)}{g(xc_1 - ry_1)} \\ D &= \frac{2zc_1 - 2c_2udy_1 - xu^2dc_1 + 2y_2udc_1 - 2\chi y_1 + ru^2dy_1}{g(xc_1 - ry_1)} \end{aligned}$$

This shows that in the plane $\Pi_{\gamma, \omega, u}$ using coordinates (s, t) , the walls are semicircles with centres $(C, 0)$ and radii $R = \sqrt{D + C^2}$.

Notation 2.1. We will generally use C and R to denote the centre and radius of any pseudo-wall in question. To make the parameters clearer we will occasionally write $C = C_v(w)$ and $R = R_v(w)$ emphasising the fact that these are functions of a variable w indexed by a parameter v .

Remark 2.2. In the case where $\mu_\omega(w) = \mu_\omega(v)$ there are no square terms and the wall is a line of the form $s = y_1/x$ which, for completeness, we can view as a circle of infinite radius.

We now want to show that the dependence of the radii on w is entirely through the centres.

Lemma 2.3. *With the notation as above, if $x \neq 0$,*

$$D = \frac{ud(2y_2 - ux) + 2z}{gx} - \frac{2y_1}{x}C.$$

Proof. Simplify $D + \frac{2y_1}{x}C$. The details are left to the reader. □

When $x = 0$ we do not have to consider the radii:

Lemma 2.4. *If $x = 0$ and $y_1 > 0$ then $C = \frac{z + duy_2}{gy_1}$. In particular, the centre is independent of w .*

We leave the elementary proof as an exercise for the reader.

Remark 2.5. So, if $x > 0$, the radius is

$$R = \sqrt{D + C^2} = \sqrt{\left(C - \frac{y_1}{x}\right)^2 - F},$$

where

$$(2.1) \quad F = \frac{d}{g} \left(u - \frac{y_2}{x}\right)^2 + \frac{1}{x^2 g} (y_1^2 g - y_2^2 d - 2xz).$$

The last term in F is $\frac{1}{x^2 g}$ times $\Delta(E) - \alpha^2$, where $\Delta(E)$ is the discriminant of E . If we assume v is a Bogomolov class then this is non-negative ($\alpha^2 \leq 0$ by the Hodge Index Theorem). Since, $d \geq 0$ by the Hodge Index Theorem we have $F \geq 0$ for all u .

2.1. Real constraints. We can now prove:

Proposition 2.6. *Suppose v is a Bogomolov class. Then in $\Pi_{\gamma, \omega, u}$, the real pseudo-walls are **nested**. By nested we mean that if W_1 and W_2 are two distinct (semi-circular) pseudo-walls in $\Pi_{\gamma, \omega, u}$ then either W_1 is entirely contained in the interior of the semi-circle W_2 or vice-versa. In particular, distinct pseudo-walls cannot intersect and one pseudo-wall cannot be entirely outside any other pseudo-wall.*

Proof. When $x = 0$ this follows immediately from Lemma 2.4 as the centres of the circles are fixed. Note that $y_1 > 0$ unless v is the class of a sheaf supported in dimension 0. So we suppose that $x \neq 0$. The result will follow from Lemma 2.3 if we can show that $C + R$ and $R - C$ are both monotonic increasing (or decreasing) as functions of C . Note that

$$\frac{d}{dC}(C \pm R) = 1 \pm \frac{1}{R} \left(C - \frac{y_1}{x}\right).$$

But $F \geq 0$ for all u by the Remark 2.5. Hence, $R \leq |C - y_1/x|$ and so if $C \neq y_1/x$, $\left|\frac{d}{dC}(C + R)\right| \geq 0$ with equality if and only if $F = 0$. Similarly $\left|\frac{d}{dC}(C - R)\right| \geq 0$ for all $C \neq y_1/x$ but has the opposite sign to the $C + R$ case. If $F = 0$ then

$$\{C + R, C - R\} = \left\{2C - \frac{y_1}{x}, \frac{y_1}{x}\right\}$$

and again we are done. \square

Note that two pseudo-walls may be coincident. In fact, there will typically be continuous families of w giving coincident pseudo-walls (by solving $C = C_0$ for w).




FIGURE 1. Typical real pseudo-walls

Remark 2.7. Now $R > 0$ exactly when

$$\left(C - \frac{y_1}{x}\right)^2 > F \geq 0.$$

So either $C > y_1/x + \sqrt{F}$ or $C < y_1/x - \sqrt{F}$. But, by Remark 1.8, we also have $y_1/x > sg$ and so $C < y_1/x$. Then $\frac{d}{dC}(C - R) \geq 2$. Note also that $R \rightarrow \infty$ as $C \rightarrow -\infty$. Consequently, we have that any point $(s, t) \in \{(a, b) : a < y_1/x, b > 0\}$ is contained in a unique real pseudo-wall (see Figure 1).

Corollary 2.8. Any pseudo-wall must intersect $s = C_0$, where

$$C_0 = \begin{cases} \frac{y_1}{x} - \sqrt{F} & \text{if } x > 0. \\ \frac{x}{z + duy_2} & \text{if } x = 0 \text{ and } y_1 > 0. \\ gy_1 & \end{cases}$$

Remark 2.9. This is particularly useful when trying to locate walls as it immediately provides us with bounds on c_1 :

$$rC_0 < c_1 < y_1 + (r - x)C_0.$$

See Example 4.1 and Proposition 4.6 below.

Remark 2.10. Using the result of Lo and Qin [LQ11] which says that for each ray there is an upper bound on the radius we can now deduce that bound is uniform in s . In the next section we will give an explicit such bound but it will also follow that the bound is not uniform in u (and we give an example in a later section to show that it can be unbounded).

3. APPLICATIONS

3.1. Bertram's Nested Wall Theorem. We can use these technical results to quickly prove:

Theorem 3.1. *Let X be a smooth complex projective surface. Pick an ample $\omega \in \text{NS}(X)$ and a real class $\beta \in \text{NS}_{\mathbb{R}}(X)$. Then for any class $v \in \text{N}(X)$ representing the Chern character of a μ -stable sheaf or a torsion sheaf on X there is a half-plane $\Pi = \Pi_{\beta, \omega, 1}$ in $\text{Stab}(X)$ which contains the basic Bridgeland stability condition $(\mathcal{A}_{\beta, \omega}, Z_{\beta, \omega})$ such that the intersection of the walls with Π are nested.*

In particular, if $\rho(X) = 1$ the walls in the space of basic stability conditions are nested.

Proof. We decompose $\beta = b\omega + \gamma$ as usual with $\gamma \cdot \omega = 0$. Then Proposition 2.6 implies that the real pseudo-walls are nested. Since the actual walls are a subset of these walls, these must be nested as well. \square

One immediate corollary is that each chamber is path connected to the $t = 0$ axis:

Corollary 3.2. *For any β and ω , the moduli space of S -equivalence classes of μ_Z -semistable objects for $Z = Z_{s\omega + u\beta, t\omega}$ is isomorphic to a moduli space of μ_Z -semistable objects for some t arbitrarily small.*

This shows that Bertram's "sliding down the wall" trick used in [MM11] always works. In the case of abelian surfaces and K3 surfaces this implies that such moduli spaces are projective varieties and represent the appropriate moduli functor (see [MM11] and [MY11b] for special cases of this).

3.2. A lower bound on the centres. We have seen in Remark 2.7 that there is an upper bound for the centres of the walls. We now show that the centres of actual walls are also bounded below as well (at least for a large class of v). This follows theoretically from the work of Lo and Qin ([LQ11, Theorem 1.1(ii)]) who show that in each ray there are no walls for t sufficiently large. Their construction is not constructive and our aim here is to use the Nested Wall Theorem to compute explicit bounds.

We prove this in a series of lemmas. For these technical lemmas we need some parameters from the surface. Let $\tilde{g} = \gcd_E(c_1(E) \cdot \omega)$. Then $g/\tilde{g} = g'$ is an integer and $y_1 g'$ is always an integer (even though y_1 may not be). For any v we let $p = \gcd(r(v), c_1(v) \cdot \omega/\tilde{g}) = \gcd(x, y_1 g')$ in the notation of the previous section. What we do in the lemma is to show that if we choose s sufficiently close to y_1/x we can ensure that there are no subsheaves of E in A_s . We have to work harder to show that there can be no μ_Z -destabilising objects of rank bigger than $r(E)$. When $p = 1$ it turns out that the same bound works but we have to go closer to y_1/x if $p > 1$.

The bound is constructed as follows. Using Euclid's algorithm, consider the set of pairs of integers (m, n) such that $g'ny_1 - mx = p$. We assume $n > 0$. Fix some (m_0, n_0) and then observe that any other pair is given by $(m, n) = (m_0 + y'\lambda, n_0 + x'\lambda)$ for some integer λ , where $y' = g'y_1/p$ and $x' = x/p$. Note that the ratio m/n tends to y'/x' as $n \rightarrow \infty$. So

given any positive real number ξ , we can pick some λ such that the set

$$\left\{ (j, k) : \frac{m + \lambda y'}{n + \lambda x'} < \frac{j}{k} < \frac{y'}{x'}, \gcd(j, k) = 1, 0 < k \leq \xi x \right\}$$

is empty. Let us, for the moment, denote the least choice of such m and n for a given ξ by m_0 and n_0 and define the set

$$P_\xi = \{(m, n) : g'ny_1 - mx = p \text{ and } m \geq m_0, n \geq n_0, n > \frac{p}{g'} \sqrt{\frac{g}{\Delta}}\},$$

where $\Delta = gy_1^2 - dy_2^2 - \alpha^2$. This last mysterious condition on n often it imposes no constraint but it will ensure that any wall which is sufficiently near y_1/x must also cross $m/g'n$ whenever $(m, n) \in P_\xi$.

Now suppose E is a torsion-free sheaf with $v(E) = v$. Then $\mu_\omega(E) = \frac{m\tilde{g}}{n} + \frac{\tilde{g}}{nx'} > 0$ and so it is reasonable to ask that $E \in \mathcal{A}_{s\omega+u\gamma, \omega}$.

Our strategy is as follows:

- (1) We show that a wall does cross $s = m/g'n$ and some other predetermined line closer to $s = y_1/x$ then the rank of any destabilizer must be at most ξx for a suitable ξ depending only on v and X . For $p = 1$ we do not require this extra assumption and we can set $\xi = 1$.
- (2) Assuming $F > 0$, we show for any $(m, n) \in P_\xi$ for the ξ given in (1) then any object which is μ_Z -stable for any Z in the region $(m/g'n, y_1/x) \times \mathbb{R}_{>0}$ must be a μ_ω -semistable sheaf.
- (3) We can then deduce that object must be μ_Z -stable for all Z in the region.
- (4) For $F = 0$ we show directly that there are no walls at all.
- (5) We also need to assume $x > 0$ but we will finish this section by showing that y_1 is always an upper bound for R when $x = 0$.

First of all, set $\xi = 1$.

Lemma 3.3. *With this notation, if $E \in \mathcal{A}_{s\omega+u\gamma, \omega} = \mathcal{A}_s$ is a sheaf then E must be μ_ω -semistable and for any t and $s \geq m_0/g'n_0$ any $Z_{s\omega+u\gamma, t\omega}$ -destabilizing subobject in \mathcal{A}_s must have rank strictly bigger than $r(E)$ or it has the same slope and μ_Z -destabilizes E for all $s < y_1/x$ and all t .*

Proof. Since $\frac{y_1 g}{x} \geq \mu_\omega^-(E) > sg > \frac{\tilde{g}m}{n}$ and if E is not μ_ω -semistable then $\mu_\omega^-(E)$ is given by $\mu(F)$ for some sheaf F with $r(F) < r(E)$, which contradicts the definition of m and n . So E must be μ_ω -semistable.

Suppose that $k \rightarrow E$ destabilizes E in \mathcal{A}_s with quotient q . Then k must be a sheaf K . Suppose $v(K) = w = (r, c_1\omega + c_2\gamma + \alpha', \chi)$. Then we have a short exact sequence $K/Q^{-1} \rightarrow E \rightarrow Q^0$. Since $c_1(Q_{\text{free}}^0) \cdot \omega \leq c_1(Q^0) \cdot \omega$ and both K/Q^{-1} and Q^0 are in T , we must have one of them with slope less than or equal to the slope of E and this is impossible unless it μ_Z -destabilizes for all $s < y_1/x$ by Remark 2.2. So Q^0 must be supported on points. Since Q^{-1} must be torsion-free, we also have that $r > x$ as required. \square

For the rank bigger than x case we appeal to a different lemma. Observe first that, for any ξ and $(m, n) \in P_\xi$ we have

$$\frac{m}{g'n} + \frac{2\tilde{g}p - 1}{2gnx} = \frac{y_1}{x} - \frac{1}{2gnx}.$$

What we do now is to show that the critical wall for a destabilizing object cannot lie outside the wall unless the rank is at most $r(v)$.

Lemma 3.4. *Pick any $\xi > 0$ and $(m, n) \in P_\xi$. Suppose $p = 1$ and $0 \rightarrow k \rightarrow e \rightarrow q \rightarrow 0$ is a short exact sequence with $v(e) = v$ and $\mu_Z(q), \mu_Z(k) < \infty$ in \mathcal{A}_s for $s = \frac{m}{g'n}$ then $r(k) \leq r(e)$ and $r(q) \leq r(e)$.*

Proof. Let $\text{ch}(k) = (r, c_1\omega + c_2\gamma + \alpha', \chi)$. Since, Z is a stability condition we have $0 < \text{Im } Z(k) < \text{Im } Z(e)$. Suppose $r(k) > r(e)$ then for $s = m/g'n$ we have

$$(3.1) \quad 0 < c_1 - r \frac{m}{g'n} < y_1 - x \frac{m}{g'n} = \frac{g'n y_1 - m x}{g'n} = \frac{p}{g'n}.$$

Clearing denominators, we have $0 < c_1 g'n - r m < p = 1$. But $c_1 g'n - r m$ is an integer. Similarly for q . \square

Lemma 3.5. *Suppose $p > 1$. Let $(m, n) \in P_\xi$ for $\xi = \frac{2\tilde{g}p}{2\tilde{g}p - 1}$ and $0 \rightarrow k \rightarrow e \rightarrow q \rightarrow 0$ is a short exact sequence with $v(e) = v$ and $\mu_Z(e), \mu_Z(k) < \infty$ in \mathcal{A}_s for both $s = \frac{m}{g'n}$ and also for $s = \frac{m}{g'n} + \frac{2\tilde{g}p - 1}{2gnx}$. Then $r(k) \leq r(e)$ and $r(q) \leq r(e)$.*

Proof. Let $\text{ch}(k) = (r, c_1\omega + c_2\gamma + \alpha', \chi)$. Since, $c_1 - r s < y_1 - x s$ and $s < y_1/x$, we have that $\mu_\omega(k) < \mu_\omega(e)$. Suppose $p > 1$ and $k \rightarrow e$ a subobject for the other value of s in the lemma then we have

$$0 < c_1 - r \left(\frac{m}{g'n} + \frac{2\tilde{g}p - 1}{2gnx} \right).$$

Rearranging, we have

$$r \frac{2\tilde{g}p - 1}{2\tilde{g}x} < c_1 g'n - m r.$$

Combining with (3.1) gives

$$r < \frac{2\tilde{g}p}{2\tilde{g}p - 1} x = \xi x.$$

But this is impossible for our choice of (m, n) . The same argument works for q . \square

The condition about infinite μ_Z -slope will not be relevant because this can only happen if the destabilizer exists for all $s < y_1/x$ and all t by Remark 2.2. We can now treat the $F = 0$ case.

Theorem 3.6. *If $F = 0$ then there are no walls in $\mathcal{A}_{s\omega + u\gamma, \omega}$ for any s and u except for $s = y_1/x$.*

Proof. When $F = 0$, we have $C + R = y_1/x$ for any wall. Consequently, for any $\xi > 0$, any such wall must intersect some $s = m/g'n$ for some $(m, n) \in P_\xi$. It must also intersect $s = \frac{y_1}{x} - \frac{1}{2g'n}$ and this contradicts our three lemmas. \square

- Notes 3.7.** (1) The condition $F = 0$ is actually rather strong. It implies that $u = y_2/x$ (or $\gamma = 0$) and the discriminant $\Delta(E) = \alpha^2$. Then the Hodge Index Theorem implies that $\alpha = 0$ and $\Delta(E) = 0$. So these are Bogomolov critical sheaves.
- (2) More generally, line bundles on any smooth projective surface always satisfy $F = 0$ and the conclusion of the Theorem is already well known (see [AB09, Prop 3.6(b)]).
- (3) When $\rho(X) = 1$ and $K_X = 0$ then $F = 0$ on a K3 surface only for line bundles but for abelian surfaces this happens exactly for semi-homogeneous bundles and we conclude that these are always μ_Z -stable for all Z . This gives a proof independently of Fourier-Mukai transform methods.
- (4) It should be possible to repeat the analysis and these theorem for the refined basic stability condition given for K3 surfaces by Bridgeland ([Bri08] and used in [ABL07] and [MY11a]) in which v is multiplied by $\sqrt{\text{Td}(X)}$. In that case the critical objects should correspond to spherical objects.

We can now assume $F > 0$. Note that if $(m, n) \in P_\xi$ for any $\xi > 0$ then we require

$$(3.2) \quad n > \frac{p}{g'} \sqrt{\frac{g}{\Delta}}.$$

We now use this condition by observing that $F > \frac{\Delta}{x^2 g} > \frac{p^2}{g'^2 n^2 x^2}$ for all u and so

$$C < \frac{y_1}{x} - \sqrt{F} < \frac{y_1}{x} - \frac{p}{g'n} = \frac{m}{g'n}.$$

So any wall which crosses any line $s = s_0 > m/g'n$ for any $(m, n) \in P_\xi$ and $\xi \geq 1$ must also cross $s = m/g'n$ as well. In particular, this holds for the least such n .

Now let $\xi_p = 1$ if $p = 1$ and $\xi_p = \frac{2gp}{2\tilde{g}p - 1}$ if $p > 1$.

Lemma 3.8. *Let $s_0 = m/g'n$ for some $(m, n) \in P_\xi$ for $\xi = \xi_p$. Suppose e is an object of some \mathcal{A}_s for all $s_0 < s < y_1/x$ with $v(e) = v$ and $r(v) > 0$. Suppose for (at least) one s there is a real number $t > 0$ such that e is μ_Z -stable with $Z = Z_{s\omega + u\gamma, t\omega}$. Then e must be a μ_ω -semistable sheaf.*

Proof. By lemmas 3.3, 3.4 and 3.5 it suffices to show that e must be a sheaf. But by lemmas 3.4 and 3.5 the rank of E^{-0} must be less than or equal to r . But then $r(E^{-1}[1]) = 0$ as it cannot be positive. But E^{-1} is torsion-free, so $E^{-1} = 0$, as required. \square

So what we have shown is that there is some $s_0 < y_1/x$ (which can be computed explicitly) with the following property. If e is an object of some \mathcal{A}_s for $s_0 < s < y_1/x$ and is μ_Z -stable for some t then it must be a μ_ω -semistable sheaf by 3.8. But then by lemmas 3.3, 3.4 and 3.5 it must be μ_Z -stable for all $s_0 < s < y_1/x$ and $t > 0$. What this means is

that no wall can intersect either the region $s \geq \frac{m}{g'n}$ (when $p = 1$) or $s \geq \frac{2gny_1 - 1}{2gnx}$ (when $p > 1$). For the moment combine these to give the bound

$$\frac{y_1}{x} - \frac{(2p\tilde{g} - 1)\epsilon + 1}{2gnx},$$

where $\epsilon = 1$ when $p = 1$ and $= 0$ otherwise. Then $C + R \leq \frac{y_1}{x} - \frac{(2p\tilde{g} - 1)\epsilon + 1}{2gnx}$. Solving for C we obtain:

Proposition 3.9. *Suppose v is the Chern character of a μ_ω -stable sheaf. Then the centres C of the walls in $\Pi_{\gamma, \omega, u}$ are bounded below by*

$$\begin{aligned} & \frac{1}{2} \left(\frac{mx + g'ny_1}{g'nx} - Fg'nx \right) \text{ when } p = 1 \text{ or} \\ & \frac{4gny_1 - 1}{4gnx} - Fg'nx \text{ when } p > 1. \end{aligned}$$

Combining this with Proposition 2.6 we have the main theorem of the paper:

Theorem 3.10. *Fix a Chern character v of a μ_ω -stable sheaf or a torsion sheaf with $F > 0$. In the half-plane $\Pi_{\beta, \omega}$ parametrized by (s, t) with $t > 0$, there are real numbers C_0 and R_0 such that all of the walls corresponding to v are contained in the semi-circle with centre $(C_0, 0)$ and radius R_0 . In particular, the radii of the walls are bounded above by R_0 .*

Notes 3.11. The bounds are not always sharp. This is especially true for $p > 1$ where the $p = 1$ bound may often suffice. But even when $p = 1$, we shall see an example below where the bound is not quite sharpest. The bounds are constructive (we shall give explicit examples in the next section). For the first three points of this note of this note we assume $p = \gcd(x, y_1g') = 1$.

(1) For $(m, n) \in P_1$ with n least

$$\frac{1}{2} \left(\frac{mx + g'ny_1}{g'nx} - Fg'nx \right) \leq C < \frac{y_1}{x} - \sqrt{F},$$

and

$$0 < R \leq \frac{1}{2} \left(Fg'nx - \frac{1}{g'nx} \right),$$

where F is given by Equation 2.1 $g' = g / \gcd_E(c_1(E) \cdot \omega)$.

(2) The upper and lower bounds on s are

$$\frac{y_1}{x} - Fg'nx \leq s < \frac{m}{g'n}.$$

(3) Recall the construction of m and n : they are given by Euclid's algorithm applied to $ng'y_1 - mx = 1$ and such that there are no integers j and $0 < k \leq x$ such that

$$\frac{m}{n} < \frac{j}{k} < \frac{y_1g'}{x}.$$

We also had the technical assumption on n given by 3.2 but this would rarely need to be imposed as the RHS is often less than 1. It also makes sense to choose m to be the least such. For example (ignoring 3.2), suppose $g' = 1$ then if $y_1 = 2$ and $x = 1$, we can pick $m = 1 = n$. Whereas, if $y_1 = 10$ and $x = 5$ then we pick $m = 7$ and $n = 4$.

- (4) The dependence on u is quadratic entirely through F . In particular, it is potentially unbounded as u increases. We shall see an example below where the walls are actually unbounded in the (u, s) -plane.

When $x = 0$ we proceed differently. Recall from Lemma 2.4 that all walls have a fixed centre:

$$C = \frac{z + duy_2}{gy_1}.$$

Observe that if $w = (r, (c_1\omega + c_2\gamma + \alpha'), \chi)$ corresponds to a wall then $r > 0$ because then $\mu_Z(w) - \mu_Z(v)$ would be constant. But using Remark 2.9, we have

$$C < \frac{c_1}{r} < \frac{y_1}{r} + C \leq y_1 + C.$$

In other words, the slope of w is bounded by constants. But, an object $e \in \mathcal{A}_s$ with $\text{ch}(e) = w$ must satisfy $\mu_\omega(e) > s$ and so if the wall has radius $R = R_v(w)$, we have $C + R < \frac{c_1}{r} < y_1 + C$ and so $R < y_1$. So we have proved:

Proposition 3.12. *If $x = 0$ then the radius of any wall for $v = (0, y_1\omega + y_2\gamma + \alpha, z)$ has radius less than y_1 .*

Combining the theorem with local finiteness of the walls we can state a version of Theorem 1.1(i) in [LQ11]:

Theorem 3.13. *Suppose $F > 0$ for some u . Then, for an $s \neq \frac{y_1}{x} - \sqrt{F}$, there are only a finite number of mini-walls in the ray $\Xi_{s\omega+\gamma, \omega}$.*

4. EXAMPLES AND COUNTEREXAMPLES

4.1. Rank 0 case.

Proposition 4.1. *Suppose (X, ω) is a surface with $\rho(X) = 1$ and $g' = 1$. Then there are no walls in the basic Bridgeland stability plane for $v = (0, 1, kg)$ for any integer k .*

Proof. Let E be a torsion sheaf with $\text{ch}(E) = v$ (a twist of the structure sheaf of a divisor in the linear system of ω) and suppose $K \rightarrow E$ is a (maximally) destabilizing object in \mathcal{A}_s for some s with $\text{ch}(K) = (r, c, \chi)$. Observe that K cannot have torsion because if it did K_{tors} would have to have $c = 1$ and so E/K_{tors} would be a skyscraper and so $\mu_Z(K) < \mu_K(K_{\text{free}})$ contradicting the maximality. So (at least) $r > 0$. It also follows by a similar argument that K is μ_ω -semistable. We use Remark 2.9. We have $C_0 = z/g = k$ and so we have

$$kr < c < 1 + kr,$$

which is a contradiction. □

When $z \neq kg$ then there may be walls. For example, consider $E = \mathcal{O}(\omega)/\mathcal{O}_X$ for some map $\mathcal{O}_X \rightarrow \mathcal{O}(\omega)$. This is just a line bundle supported on a divisor in the linear system $|\omega|$. This has Chern character $(0, \omega, g/2)$ and there is a wall with centre $1/2$ and radius $1/2$ given by $K = \mathcal{O}(\omega)$.

4.2. Rank 1 case. We now consider the cases studied in [MM11]. There $X = (\mathbb{T}, \ell)$ was an irreducible principally polarized abelian surface (ppas for short) and so $\rho(X) = 1$ and $g = 2 = \tilde{g}$. Note that we must have $\gamma = 0$ (and so $d = 0$ in the formulae of Note 3.11). We set $v = (1, 2\ell, 4 - k)$ and $\beta = \omega$. Then $F = k$ and the bounds become

$$\frac{3 - k}{2} \leq C < 2 - \sqrt{k},$$


$$R \leq \frac{1}{2}(k - 1),$$

and the walls are constrained to satisfy $2 - k \leq s < 1$. It follows that $L^2\mathcal{I}_p$ for a point p is σ -stable for all $\sigma \in \Pi_{\omega, \omega, 0}$ for which the sheaf is in \mathcal{A} . Note that, in that case, the maximal radius is achieved (by $w = (1, \ell, 1)$). This agrees with the finding of [MM11].

In that paper (and elsewhere) it is shown that the twisted ideal sheaf $L \otimes \mathcal{I}_V$ is Z -stable for any zero-scheme V but the situation is very different if we assume \mathbb{T} is reducible. Suppose $\mathbb{T} = E_1 \times E_2$ is a product of elliptic curves (for simplicity, we assume they are non-isomorphic but this is not really necessary). Then the canonical polarization splits as $\ell = \omega = \ell_1 + \ell_2$. In this case, we have $\tilde{g} = 1$ and so $g' = 2$. This time we must pick $m = 3$ and $n = 1$. We still have $F = k$ but now $s < 3/2$ for our walls and our maximal radius (of a pseudo-wall) is now $k - 1/4$ with centre $7/4 - k$. But this does not correspond to a real wall. The maximal wall is given by $w = (1, 2\ell_1 + \ell_2, 2)$ with centre $2 - k$ and radius $\sqrt{k(k - 1)}$. Then $s < 2 - k + \sqrt{k(k - 1)} < 3/2$, for all k . In fact, the limit of $C + R$ for these walls as $k \rightarrow \infty$ is $3/2$ but the limit is never achieved.

To give a more complete picture, consider the same product torus but this time let $v = (1, \ell_1 + \ell_2, 1 - k)$. This corresponds to the sheaves $L \otimes \mathcal{I}_V \otimes \mathcal{P}_{\hat{x}}$, where $\mathcal{P}_{\hat{x}} \in \text{Pic}^0 \mathbb{T}$ and $|V| = k$. These sheaves have two families of potentially destabilising subsheaves: $L_1 \otimes \mathcal{I}_{V'}$ and $L_2 \otimes \mathcal{I}_{V'}$ up to twists by flat line bundles, where $|V'| < |V|$. These correspond geometrically to $V' \subset V$ and $V \setminus V'$ being contained in a translate of either E_2 or E_1 , respectively. These two families of walls intersect along plane of constant u (as predicted by Theorem 3.1). The situation is illustrated in Figure 2 for the case $k = 4$, where the vertical axis is t and the two grey planes are $s = 1$ and $u = 0$. On the plane $u = 0$ the walls have centres at $s = m - 3$ and radii $\sqrt{m^2 - 8m + 12}$. These radii are positive for $m = 0$ and $m = 1$ only. Note that in the (s, u) plane the walls are parabolas and so are unbounded.

4.3. Global finiteness. We can ask if the set of walls is always finite but two examples in [Mea12] show that this is not the case. For an irreducible principally polarized abelian surface (\mathbb{T}, ω) the Chern characters $(1, 2\omega, 2)$ and $(1, 2\omega, 1)$ result in infinite families of walls. But it does confirm our finding that, away from C_0 , each ray contains only finitely many walls. However, we can prove:

FIGURE 2. Walls in $\Pi_{\ell_1, \ell, u}$

Proposition 4.2. *In the case of a principally polarized abelian surface (\mathbb{T}, ω) , if q is rational then the number of mini-walls in the ray $\beta = q\omega$ is finite.*

Proof. This follows because for such q , there is a Fourier-Mukai transform Φ_q which reverses the ray (see [MM11, Prop 3.2]). Then the proposition follows from Theorem 3.10. \square

As was shown by Bridgeland in [Bri08], the space of all (good) Bridgeland stability conditions $\text{Stab}(\mathbb{T})$ is 3-dimensional and each stability condition can be written as $\Phi(\mathcal{A}, Z)$ for some Fourier-Mukai transform Φ . Since Φ preserves the chamber and wall structure we have

Theorem 4.3. *Let (\mathbb{T}, ω) be an irreducible principally polarized abelian surface. The walls in $\text{Stab}(\mathbb{T})$ are nested real hypersurfaces.*

Remark 4.4. This will hold more generally for abelian surfaces and K3 surfaces when stated properly but we omit the details.

Example 4.5. In particular, by Theorem 3.13, if $y_1/x - \sqrt{F}$ is rational (ie, when F is a square rational) then the number of walls in $\text{Stab}(X)$ is (globally) finite. For example, on a ppas (\mathbb{T}, ω) with $v = (5, 3\omega, 1)$ we have $F = 4/25$ and then the critical ray is $s = 1/5$.

4.4. The case of O’Grady’s Holomorphic Symplectic Space. Let us now look at an example where $p \neq 1$. In [O’G03], Kieran O’Grady showed that the moduli space of Gieseker stable sheaves with Chern character $(2, 0, -2)$ (modulo natural torus actions) gives a simply connected symplectic holomorphic manifold which was not one of the (then) known types.

Proposition 4.6. *Let (X, ω) be an irreducible principally polarized abelian surface. Then the Chern character $v = (2, 2\omega, 0)$ has no walls in $\text{Stab}(X)$.*

Proof. Since $\text{Stab}(X)$ is generated by Fourier-Mukai transforms it suffices to prove this for basic Bridgeland stability conditions. It suffices to show that any μ_ω -semistable sheaf E which is also μ_Z -stable for some Z is μ_Z -stable for all Z . Note that $F = 1$ for this choice of v and $C_0 = 0$. So Remark 2.9 gives $0 < c < 2$ and then $c_1(K) = \omega$ for any destabilizing object K in any \mathcal{A}_s . We also must have $r(K) > 1$ because otherwise $xc_1 - y_1r = 0$. As usual, we can assume that K is μ_ω -semistable. On the other hand, the centre of the wall corresponding to (r, ω, k) is $\frac{-\chi}{2r-2} < C_0 = 0$. But then $\chi > 0$ and that contradicts the Bogomolov inequality for K . \square

4.5. A Conjecture. The examples above and Theorem 3.1 and its corollary suggest the following:

Conjecture 4.7. For any smooth complex projective surface X and any $v \in N(X)$ the complement of the walls in the space of good stability conditions union the boundary of its closure is path-connected.

In other words, any σ -stable moduli spaces survives to the boundary of $\text{Stab}(X)$. The conjecture holds for abelian surfaces and holds for the subspace of basic stability conditions for any surface.

REFERENCES

- [AB09] Daniele Arcara and Aaron Bertram. Reider’s theorem and Thaddeus pairs revisited. arxiv:09043500, 2009.
- [ABL07] Daniele Arcara, Aaron Bertram, and Max Lieblich. Bridgeland-stable moduli spaces for K-trivial surfaces. arxiv: 07082247, to appear J. European Mathematical Society, 2007.
- [Bri07] Tom Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.
- [Bri08] Tom Bridgeland. Stability conditions on K3 surfaces. *Duke Math. J.*, 141(2):241–291, 2008.
- [Kaw11] Kotaro Kawatani. Stability of Gieseker stable sheaves on K3 surfaces in the sense of Bridgeland and some applications. arxiv:1103.3921, 2011.
- [LQ11] Jason Lo and Zhenbo Qin. Mini-walls for Bridgeland stability conditions on the derived category of sheaves over surfaces. arxiv:1103.4352, 2011.
- [Mea12] Ciaran Meachan. Phd thesis. forthcoming, 2012.
- [MM11] Antony Maciocia and Ciaran Meachan. Rank one Bridgeland stable moduli spaces on a principally polarized abelian surface. arxiv:1107.5304, to appear Int. Math. Research Notices, 2011.
- [MY11a] Hiroki Minamide, Shintarou Yanagida, and Kota Yoshioka. Fourier-Mukai transforms and the wall-crossing behavior for Bridgeland’s stability conditions. arXiv:1106.5217, June 2011.

- [MY11b] Hiroki Minamide, Shintarou Yanagida, and Kota Yoshioka. Some moduli spaces of Bridgeland's stability conditions. arXiv:1111.6187, November 2011.
- [OG03] Kieran G. O'Grady. A new six-dimensional irreducible symplectic variety. *J. Algebraic Geom.*, 12(3):435–505, 2003.

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF EDINBURGH, THE KING'S BUILDINGS, MAYFIELD ROAD, EDINBURGH, EH9 3JZ.,
E-mail address: A.Maciocia@ed.ac.uk